YMCA UNIVERSITY OF SCIENCE& TECHNOLOGY, FARIDABAD

B. TECH. 3RD SEMESTER (UNDER CBCS)

NUMERICAL METHODS (ME-211C) Max. Marks:75 Time: 3 Hours Note: 1. It is compulsory to answer the questions of Part -1. 2. Answer any four questions from Part -2 in detail. 3. Different parts of the same question are to be attempted adjacent to each other. 4. Assume suitable standard data wherever required, if not given. PART-1 (1.5)Q1 (a) Why Secant method is used (1.5)(b) What are the limitations of Gauss elimination method over Jordon method? (1.5)(c) Define Interpolation. (1.5)(d) Write Weddle rule formula. (1.5)(e) What do you mean by Least square method? (1.5)(f) What are the applications of numerical methods in industries? (1.5)(g) What do you mean by 5-point standard formula? (1.5)(h) Why corrector is used in Miline's method. (1.5)(i) Define error and accuracy. (1.5)What do you mean by difference operators. PART-2 Q2 a) Using the Regula Falsi method, compute the real root of the equation (7) $xe^{x}=2$, in (0.8,0.9) correct to four decimal places. (8) Solve by Jacobi iteration method, correct to three decimal places 10x+y-z=11.19x+10y+z=28.08-x+y=10z=35.61Define interpolation and Estimate the value of f(22) and f(42) from the following Q3 available data: (15)45 X: 20 35 40 204 F(X): 354 332 291 260 231 A jet fighter's position on an aircraft carrier's runway was timed during landing. Q4 1.5 1.6 1.2 1.3 1.4 t, (sec.) 1.0 1.1 y, (m) 7.989 8.403 8.781 9.129 9.451 9.750 10.031 (15)Where y is the distance from the end of carrier. Estimate velocity dy/dt and acceleration d^2y/d^2t at (i) t = 1.1, (ii) t = 1.6 using numerical differentiation.

Given $y' = xy + y^2$, y(0) = 1, find y(0.1) by Eulers's method, y(0.2) by Euler's 05 modified method, y(0.3) by Runga Kutta Method order 4th and y(0.4) by Adam's (15) method.

Solve the Laplace equations using 5 point standard and Diagonal formula Q6 (15)

(15)What is crank Nicolson method? Derive the equation. Q7.