



### STANDARD NORMAL TABLE (Z)

Entries in the table give the area under the curve between the mean and z standard deviations above the mean. For example, for  $z = 1.25$  the area under the curve between the mean (0) and z is 0.3944.

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.0000	0.0040	0.0080	0.0120	0.0160	0.0190	0.0239	0.0279	0.0319	0.0359
0.1	0.0398	0.0438	0.0478	0.0517	0.0557	0.0596	0.0636	0.0675	0.0714	0.0753
0.2	0.0793	0.0832	0.0871	0.0910	0.0948	0.0987	0.1026	0.1064	0.1103	0.1141
0.3	0.1179	0.1217	0.1255	0.1293	0.1331	0.1368	0.1406	0.1443	0.1480	0.1517
0.4	0.1554	0.1591	0.1628	0.1664	0.1700	0.1736	0.1772	0.1808	0.1844	0.1879
0.5	0.1915	0.1950	0.1985	0.2019	0.2054	0.2088	0.2123	0.2157	0.2190	0.2224
0.6	0.2257	0.2291	0.2324	0.2357	0.2389	0.2422	0.2454	0.2486	0.2517	0.2549
0.7	0.2580	0.2611	0.2642	0.2673	0.2704	0.2734	0.2764	0.2794	0.2823	0.2852
0.8	0.2881	0.2910	0.2939	0.2969	0.2995	0.3023	0.3051	0.3078	0.3106	0.3133
0.9	0.3159	0.3186	0.3212	0.3238	0.3264	0.3289	0.3315	0.3340	0.3365	0.3389
1.0	0.3413	0.3438	0.3461	0.3485	0.3508	0.3531	0.3554	0.3577	0.3529	0.3621
1.1	0.3643	0.3665	0.3686	0.3708	0.3729	0.3749	0.3770	0.3790	0.3810	0.3830
1.2	0.3849	0.3869	0.3888	0.3907	0.3925	0.3944	0.3962	0.3980	0.3997	0.4015
1.3	0.4032	0.4049	0.4066	0.4082	0.4099	0.4115	0.4131	0.4147	0.4162	0.4177
1.4	0.4192	0.4207	0.4222	0.4236	0.4251	0.4265	0.4279	0.4292	0.4306	0.4319
1.5	0.4332	0.4345	0.4357	0.4370	0.4382	0.4394	0.4406	0.4418	0.4429	0.4441
1.6	0.4452	0.4463	0.4474	0.4484	0.4495	0.4505	0.4515	0.4525	0.4535	0.4545
1.7	0.4554	0.4564	0.4573	0.4582	0.4591	0.4599	0.4608	0.4616	0.4625	0.4633
1.8	0.4641	0.4649	0.4656	0.4664	0.4671	0.4678	0.4686	0.4693	0.4699	0.4706
1.9	0.4713	0.4719	0.4726	0.4732	0.4738	0.4744	0.4750	0.4756	0.4761	0.4767
2.0	0.4772	0.4778	0.4783	0.4788	0.4793	0.4798	0.4803	0.4808	0.4812	0.4817
2.1	0.4821	0.4826	0.4830	0.4834	0.4838	0.4842	0.4846	0.4850	0.4854	0.4857
2.2	0.4861	0.4864	0.4868	0.4871	0.4875	0.4878	0.4881	0.4884	0.4887	0.4890
2.3	0.4893	0.4896	0.4898	0.4901	0.4904	0.4906	0.4909	0.4911	0.4913	0.4916
2.4	0.4918	0.4920	0.4922	0.4925	0.4927	0.4929	0.4931	0.4932	0.4934	0.4936
2.5	0.4938	0.4940	0.4941	0.4943	0.4945	0.4946	0.4948	0.4949	0.4951	0.4952
2.6	0.4953	0.4955	0.4956	0.4957	0.4959	0.4960	0.4961	0.4962	0.4963	0.4964
2.7	0.4965	0.4966	0.4967	0.4968	0.4969	0.4970	0.4971	0.4972	0.4973	0.4974
2.8	0.4974	0.4975	0.4976	0.4977	0.4977	0.4978	0.4979	0.4979	0.4980	0.4981
2.9	0.4981	0.4982	0.4982	0.4983	0.4984	0.4984	0.4985	0.4985	0.4986	0.4986
3.0	0.4987	0.4987	0.4987	0.4988	0.4988	0.4989	0.4989	0.4989	0.4990	0.4990
3.1	0.4990	0.4991	0.4991	0.4991	0.4992	0.4992	0.4992	0.4992	0.4993	0.4993
3.2	0.4993	0.4993	0.4994	0.4994	0.4994	0.4994	0.4994	0.4995	0.4995	0.4995
3.3	0.4995	0.4995	0.4995	0.4996	0.4996	0.4996	0.4996	0.4996	0.4996	0.4997
3.4	0.4997	0.4997	0.4997	0.4997	0.4997	0.4997	0.4997	0.4997	0.4997	0.4998

Roll No. ....

Total Pages : 4

**504101**

Mar. 2022

M.Tech. (CSE) I SEMESTER

Mathematical Foundation of Computer Science  
(MCS-18-101)

Time : 90 Minutes]

[Max. Marks : 25

Instructions :

1. It is compulsory to answer all the questions (1 mark each) of Part-A in short.
2. Answer any three questions from Part-B in detail.
3. Different sub-parts of a question are to be attempted adjacent to each other.
4. Normal distribution table is provided

### PART-A

1. (a) A biased coin has the probability of getting tail as 0.8. What is the probability of getting 4 heads in 5 trials? (1)
- (b) State Markov's inequality. (1)
- (c) State central limit theorem. (1)
- (d) What is convenience sampling? (1)

- (c) If the sample median is chosen to estimate population mean, which property of the estimator is violated? (1)
- (f) State the values of bias and variance in Goodfit, underfit and overfit models. (1)
- (g) What is the chromatic number of following graphs: K5 graph, bipartite graph, a totally unconnected graph (empty graph) of n vertices? (1)
- (h) In how many ways a team of 7 students consisting of 4 boys and 3 girls can be chosen from a group of 10 boys and 9 girls. (1)
- (i) Define the concept of membership and support in the context of fuzzy sets. (1)
- (j) Differentiate between classification and regression. (1)

### PART-B

2. (a) 1000 tubes lights with a mean life of 120 days are installed in a new factory. Length of their life is normally distributed with standard deviation of 20 days.
- (i) How many tube lights will expire in less than 90 days.
- (ii) How many tube lights survive beyond 150 days. (2)

- (b) On the average one in 400 items is defective. If the items are packed in a box of 100, what is the probability that any given box will contain
- (i) no defective
- (ii) less than 2 defectives (3)
3. (a) Describe systematic sampling and multistage sampling. (2)
- (b) Write a short note on Principal Component Analysis. (3)
4. (a) In how many different ways can the letters of the word 'MATHEMATICS' be arranged so that the vowels always come together? (2)
- (b) Describe the following with the help of suitable examples: Isomorphism, Eulerian path & Circuit. (3)
5. (a) Differentiate between supervised and unsupervised learning with the help of suitable example. (3)
- (b) Write a short note on data mining. (2)
6. (a) Describe the various stages of Convolutional Neural Networks in the context of Computer Vision. (2)
- (b) Use dynamic programming to match the following sequences :  
REFED and TREFKD  
Given match score = +1 mismatch penalty = 0  
Gap Penalty=-2 (3)